



University of Kentucky
UKnowledge

Kentucky Water Resources Annual Symposium

2014 Kentucky Water Resources Annual
Symposium

Mar 10th, 3:00 PM

Session 2C

Kentucky Water Resources Research Institute, University of Kentucky

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/kwrri_proceedings

 Part of the [Engineering Commons](#), [Life Sciences Commons](#), and the [Physical Sciences and Mathematics Commons](#)

Kentucky Water Resources Research Institute, University of Kentucky, "Session 2C" (2014). *Kentucky Water Resources Annual Symposium*. 8.

https://uknowledge.uky.edu/kwrri_proceedings/2014/session/8

This Presentation is brought to you for free and open access by the Kentucky Water Resources Research Institute at UKnowledge. It has been accepted for inclusion in Kentucky Water Resources Annual Symposium by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

RESULTS OF THE STATEWIDE GROUNDWATER PATHOGEN STUDY

Susan L. Mallette, PG
Kentucky Division of Water
200 Fair Oaks Lane, Frankfort, KY 40601
(502) 564-3410
susan.mallette@ky.gov

The Kentucky Division of Water (KDOW) conducted a statewide study of pathogens in groundwater as part of a nonpoint source assessment funded in part by the Clean Water Act Section 319(h). The goal of the study was to determine if domestic water sources in Kentucky, which are not regulated by the Safe Drinking Water Act (SDWA), are sources of safe drinking water to the segment of the population whose drinking water is not supplied by public water systems, and to inform private water owners of the condition of their water with respect to bacteria. Since the SDWA does not regulate private drinking water sources, this project was an opportunity for KDOW to evaluate whether Kentucky's private domestic water source users are being exposed to pathogens in their drinking water. It was also a chance for interested groundwater users to obtain water quality data on their well and/or spring water.

Two hundred and ten domestic water wells and springs were sampled throughout the state, with priority on those used as drinking water sources. Samples were analyzed for total coliform, *E. coli*, Iron Related, Sulfate Reducing, and Slime-Forming bacteria (using BART[®] kits), caffeine and the caffeine byproduct 1,7-Dimethylxanthine. Although many factors can influence the presence of bacteria in water wells, in most cases it appeared to be the construction and maintenance of the water source that had the strongest influence. Further, it was anticipated that positive *E. coli* results, coupled with positive caffeine results, would indicate direct impacts to groundwater from human waste disposal activities; six wells had detections of both parameters, but the sources of each is not yet determined.

A secondary goal of this study was outreach and education to citizens that are using private groundwater sources as drinking water supplies. All participants were informed about the importance of routine water well maintenance. Copies of KDOW's Water Well Owner's Guide and generic Groundwater Protection Plans were handed out at each site. Participants also received the results of the samples collected from their wells/springs along with a letter of explanation regarding the quality of their drinking water.

NOTES

[illegible]

CHANGES IN NUTRIENTS AND *E. coli* DURING TWO STORM EVENTS IN
HINKSTON CREEK, KY

Tiffany L. Hunter

Brian C. Reeder

Institute for Regional Analysis and Public Policy

School of Public Affairs

Morehead State University, Morehead, KY 40351

(606) 783-2945

b.reeder@moreheadstate.edu

During storm events, rapid changes can occur in stream water nutrient loading. Hinkston Creek, KY, water quality has been analyzed for the past five years; this information was used to implement a number of best management practices (BMPs). Global warming is increasing storm intensity. More intense rainfall events (heavier rainfall over shorter periods of time) results in rapid increases in surface runoff and channel discharge; which can greatly change total pollutant loads. Accordingly, we examined short-term changes in nutrients and *E. coli* during two storm events during November and December of 2013. Water samples were taken at intervals (roughly every four hours) over the course of a few days. We took field measurements of pH, conductivity, dissolved oxygen (DO), and temperature. Water samples were brought back to the lab and immediately analyzed for *E. coli*, total suspended solids (TSS), nitrate ($\text{NO}_3\text{-N}$), soluble reactive phosphate (SRP), ammonia ($\text{NH}_4\text{-N}$), iron, sulfate (SO_4), total N (TN), and total P (TP). Sediment and sediment associated pollutants increased to a maximum just before peak discharge. These constituents remained at concentrations far above their base flow concentrations throughout the storms. Some dissolved constituents (alkalinity, and conductivity) decreased dramatically during the storms. Nitrate did not necessarily follow this trend for dissolved nutrients. During the most intense storm, nitrate concentrations increased from 700 mg m^{-3} before the storm (discharge = 400 cfs) to $>3,000 \text{ mg m}^{-3}$ at peak discharge (about 2,500 cfs). *E. coli* contamination was about

100 cfu 100 ml⁻¹ at the lowest discharge (7 cfs), which increased to over 50,000 cfu 100 ml⁻¹ at the beginning of the storm events. *E. coli* was consistently well over 1,000 cfu 100 ml⁻¹ during the storm. A couple of large storms may contribute significantly to annual nutrient loading from Hinkston Creek. Storm-event studies are vital to determining how regional nutrient loading contributes to Gulf of Mexico anoxia. Management techniques that reduce the speed of surface run-off during storms will become more important as rainfall events increase in intensity.

ENGAGING PARTNERS IN THE CANE RUN WATERSHED

Carol Hanley
Carmen Agouridis
Amanda Gumbert
Kelly Taylor
College of Agriculture, Food and Environment
University of Kentucky
859-257-3785, 257-1993
chanley@uky.edu

The Cane Run Watershed (CRW) project management team, composed of the University of Kentucky, College of Agriculture, Environmental and Natural Resources Initiative, and the Department of Biosystems and Agricultural Engineering staff, is involved in an innovative, comprehensive program that engages K-12 students and teachers and citizens within the watershed and is designed to increase understanding of water quality issues and promote urban stream restoration. The two goals of the project, one for each major audience, include the following:

Goal 1: Engage K-12 and community partners in an urban watershed, the Cane Run Watershed, in an innovative education project to improve water quality in the watershed

Goal 2: Engage all citizens within the watershed in the urban-stream restoration project

The K-12 portion of the program involves the formation of partnerships between community groups and three schools to develop a geographic education tool – a map of Cane Run Environment & Watershed. The map uses a GIS platform and is accessible from a College of Agriculture, Food and Environment web site.

At last year's symposium, the leadership team presented general features of the program and reviewed progress. At the 2014 symposium, the team will present progress on both goals, including results of the 2012-13 pre-/post-test analyses, samples of student art work and updates of the map. Upcoming plans for community involvement also will be presented. Teachers and students involved in the program will be invited to the Symposium to present their findings and talk about their maps.

NOTES

[illegible]

THE UNIVERSITY OF KENTUCKY
ENVIRONMENTAL RESEARCH AND TRAINING LAB
IS AVAILABLE FOR YOUR USE

Tricia Coakley
University of Kentucky, ERTL
161 Raymond Building, Lexington, KY 40506-0281
859-257-6757
tricia.coakley@uky.edu

The Environmental Research and Training Lab (ERTL) is a shared use core facility at the University of Kentucky offering a wide array of scientific instrumentation and methods to faculty, students, businesses, government organizations and non-profit organizations to generate the scientific data needed to achieve their research goals. We provide access to state of the art analytical equipment and customized project specific training in organic chemistry, inorganic chemistry, microbiology, and molecular biology methods.

Some of the instruments available for your use at ERTL are:

- ICP-OES, Graphite Furnace AA, and Direct Mercury Analyzer for metals
- Ion Chromatograph for nutrients and anions
- GC, GC-MS, HPLC, and LC-MS/MS for analysis of organic compounds
- Real Time PCR for gene expression assays or the quantitative analysis of DNA or RNA
- IDEXX system for the analysis of E.coli and total coliforms in water

What you will gain from conducting your research at ERTL:

- Expertise in experimental design and method development
- Well maintained scientific instrumentation covering a broad array of research areas
- A non-restrictive and supportive research environment
- Networking opportunities between academic researchers at UK, other state institutions, and industrial partners.

Researchers from many fields find a welcoming interdisciplinary environment at ERTL. Contact us to find out how we can support your research. <http://ertl.uky.edu>

NOTES

[illegible]